








POTATO

Crop Information Sheet



SUREWAY

POTATOES		Land Prep Seed Treatment	Planting In Furrow	Growth Stage 1 Sprout Development	Growth stage 2 Vegetative growth	Growth Stage 3 Tuber Initiation	Growth Stage 4 Tuber Bulking	Growth Stage 5 Maturity
Product	Description							
MicroCal	Soil pH is crucial for nutrient availability	As per soil pH, apply up to 300kg/ha.	As per soil pH, apply infurrow, up to 75kg/ha					
MicroDol								
MicroGyp	Calcium is important for aeration of soils and nutrient uptake. Calcium is also important to cell development.	Soil Amelioration - rate exceeds 60kg/ha	Fertilization - Apply in furrow at 50-60kg/ha			Fertilization - broadcast at 60 to 100kg/ha		
Rootsure	Combination product: Liquid multi nutrient and Kelp, as well as Mycorrhizae and Trichoderma	Apply to tubers before or at planting						
Tricure	For the prevention of Rhizoctonia, Stem Canker and Black Scurf		Coating - 250g - 750g per hectare. 25g TriCure, mixed with 100l of water per 100kg of seed. Applied immediately after planting, over the potatoes in furrow.	Foliar - apply at 125g to 250g/ha	Foliar - apply at 125g/ha			
Rappid	Efficient and effective liquid Polyphosphate, providing a sustainable release of available phosphate.		Drench - Apply 5L/ha		Foliar - Apply 5L/ha			
EcoKsil	Fortify plants cells and surfaces against pathogens, and increase chlorophyll concentration in the leaf.			Foliar - Apply at 2L/ha, repeat after 2 weeks				
Vitazyme	Stimulates multiple enzyme reactions, improving photosynthesis and other plant processes				Foliar - Apply at 1L/ha		Foliar - Apply 1L/ha	
Rappid K	Provides the energy required to assist with the immune system boost.					Foliar - Apply 2-5L/ha, repeat after 2-3 weeks		
Zalexin	Salicylic Acid and Multiple nutrients for foliar application during times of rapid growth and stress.					Foliar - Apply 1-2L/ha, repeat after 2-3 weeks		
Leafguard	Inhibits the infection and colonization of fungal pathogens (Late blight).					Foliar - Apply at 400ml/ha	Foliar - Apply at 400ml/ha	Foliar - Apply at 400ml/ha